

Virus isolation/detection from aseptic meningitis cases, 2008 - 2012

(Infectious Agents Surveillance Report: Data based on the reports received before November 11, 2017 from public health institutes)

2012		
1 Echovirus 6	77	21.4%
2 Coxsackievirus B5	41	11.4%
3 Echovirus 7	41	11.4%
4 Coxsackievirus A9	34	9.4%
5 Echovirus 9	27	7.5%
6 Enterovirus NT	20	5.6%
7 Rhinovirus	14	3.9%
8 Enterovirus 71	9	2.5%
9 Coxsackievirus B4	7	1.9%
10 Other enterovirus	21	5.8%
11 Mumps virus	36	10.0%
12 Other virus	33	9.2%
Total	360	100.0%

2011		
1 Echovirus 6	128	23.7%
2 Coxsackievirus B1	82	15.2%
3 Echovirus 9	51	9.4%
4 Coxsackievirus B5	50	9.3%
5 Enterovirus NT	33	6.1%
6 Coxsackievirus B4	21	3.9%
7 Echovirus 7	17	3.1%
8 Rhinovirus	14	2.6%
9 Coxsackievirus A6	7	1.3%
9 Coxsackievirus B2	7	1.3%
11 Other enterovirus	33	6.1%
12 Mumps virus	57	10.6%
13 Other virus	40	7.4%
Total	540	100.0%

2010		
1 Enterovirus 71	61	17.0%
2 Echovirus 6	39	10.9%
3 Coxsackievirus B2	25	7.0%
4 Echovirus 25	25	7.0%
5 Coxsackievirus B4	23	6.4%
6 Coxsackievirus B1	18	5.0%
7 Rhinovirus	12	3.4%
8 Enterovirus NT	11	3.1%
9 Echovirus 30	6	1.7%
10 Other enterovirus	26	7.3%
11 Mumps virus	82	22.9%
12 Other virus	30	8.4%
Total	358	100.0%

2009		
1 Coxsackievirus B3	19	8.1%
2 Enterovirus NT	17	7.3%
3 Echovirus 6	17	7.3%
4 Coxsackievirus A9	13	5.6%
5 Coxsackievirus B4	13	5.6%
6 Echovirus 30	13	5.6%
7 Echovirus 11	12	5.1%
8 Coxsackievirus B2	8	3.4%
9 Coxsackievirus B1	5	2.1%
10 Other enterovirus	29	12.4%
11 Mumps virus	45	19.2%
12 Other virus	43	18.4%
Total	234	100.0%

2008		
1 Echovirus 30	126	30.6%
2 Coxsackievirus B5	65	15.8%
3 Coxsackievirus B3	33	8.0%
4 Enterovirus NT	16	3.9%
5 Echovirus 6	13	3.2%
6 Echovirus 18	13	3.2%
7 Coxsackievirus B4	12	2.9%
8 Echovirus 9	12	2.9%
9 Coxsackievirus B1	10	2.4%
10 Other enterovirus	46	11.2%
11 Mumps virus	29	7.0%
12 Other virus	37	9.0%
Total	412	100.0%