

#### 4. Department of Bacteriology I

- 1) Iguchi A, Iyoda S, Kikuchi T, Ogura Y, Katsura K, Ohnishi M, Hayashi T, Thomson NR. A complete view of the genetic diversity of the *Escherichia coli* O-antigen biosynthesis gene cluster. DNA Research. 2015, 22(1): 101-107.
- 2) Iyoda S, Manning SD, Seto K, Kimata K, Isobe J, Etoh Y, Ichihara S, Migita Y, Ogata K, Honda M, Kubota T, Kawano K, Matsumoto K, Kudaka J, Asai N, Yabata J, Tominaga K, Terajima J, Morita-Ishihara T, Izumiya H, Ogura Y, Saitoh T, Iguchi A, Kobayashi H, Hara-Kudo Y, Ohnishi M, and EHEC working group in Japan. Phylogenetic clades 6 and 8 of enterohemorrhagic *Escherichia coli* O157:H7 with particular *stx* subtypes are more frequently found in isolates from hemolytic uremic syndrome patients than from asymptomatic carriers. Open Forum Infectious Diseases. 2014, 1(2): ofu061.
- 3) Yahata Y, Misaki T, Ishida Y, Nagira M, Watahiki M, Isobe J, Terajima J, Iyoda S, Mitobe J, Ohnishi M, Sata T, Taniguchi K, Tada Y, Okabe N. Epidemiological analysis of a large enterohaemorrhagic *Escherichia coli* O111 outbreak in Japan associated with haemolytic uraemic syndrome and acute encephalopathy. Epidemiol Infect. 2015, Jan 20:1-12.
- 4) Maeda E, Murakami K, Etoh Y, Onozuka D, Sera N, Asoshima N, Honda M, Narimatsu H, Iyoda S, Watahiki M, Fujimoto S. Does sequence type 33 of Shiga toxin-producing *Escherichia coli* O91 cause only mild symptoms? J Clin Microbiol. 2015, 53(1): 362-364.
- 5) Watahiki M, Isobe J, Kimata K, Shima T, Kanatani J, Shimizu M, Nagata A, Kawakami K, Yamada M, Izumiya H, Iyoda S, Morita-Ishihara T, Mitobe J, Terajima J, Ohnishi M, Sata T. J Clin Microbiol. 2014, 52(8): 2757-2263.
- 6) Lee K, Kusumoto M, Sekizuka T, Kuroda M, Uchida I, Iwata T, Okamoto S, Yabe K, Inaoka T, Akiba M. 2015. Extensive amplification of GI-VII-6, a multidrug resistance genomic island of *Salmonella enterica* serovar Typhimurium, increases resistance to extended-spectrum cephalosporins. Front. Microbiol. 2015, 6:78.
- 7) Matsumoto Y, Izumiya H, Sekizuka T, Kuroda M, Ohnishi M. Characterization of *blaTEM-52*-carrying plasmids of extended-spectrum-β-lactamase-producing *Salmonella enterica* isolates from chicken meat with a common supplier in Japan. Antimicrob Agents Chemother. 2014 Dec;58(12):7545-7.
- 8) Albert MJ, Al Obaid K, Alfouzan W, Sheikh AR, Udo E, Izumiya H, Bulach DM, Seemann T. Isolation of *Salmonella enterica* serovar Kentucky strain ST 198 and its H2S-negative variant from a patient: implications for diagnosis. J Clin Microbiol. 2014 Nov;52(11):4090-3.
- 9) Ido N, Lee K, Iwabuchi K, Izumiya H, Uchida I, Kusumoto M, Iwata T, Ohnishi M, Akiba M. Characteristics of *Salmonella enterica* serovar 4,[5],12:i:- as a monophasic variant of serovar Typhimurium. PLoS One. 2014 Aug 5;9(8):e104380.
- 10) Fukui D, Takahashi K, Kubo M, Une Y, Kato Y, Izumiya H, Teraoka H, Asakawa M, Yanagida K, Bando G. Mass mortality of Eurasian Tree Sparrows (*Passer montanus*) from *Salmonella* Typhimurium DT40 in Japan, winter 2008-09. J Wildl Dis. 2014 Jul;50(3):484-95.
- 11) Kobayashi T, Hayakawa K, Mawatari M, Mezaki K, Takeshita N, Kutsuna S, Fujiya Y, Kanagawa S, Ohmagari N, Kato Y, Morita M. Case report: failure under azithromycin treatment in a case of bacteremia due to *Salmonella enterica* Paratyphi A. BMC Infectious Diseases. 2014. 14:404
- 12) Ikebe T, Tominaga K, Shima T, Okuno R, Kubota H, Ogata K, Chiba K, Katsukawa C, Ohya H, Tada Y, Okabe N, Watanabe H, Ogawa M, Ohnishi M, Working Group for Beta-hemolytic Streptococci in Japan. Increased prevalence of group A streptococcus isolates in streptococcal toxic shock syndrome cases in Japan from 2010–2012. Epidemiology and Infection. 2015. 143 (4): 864-872.
- 13) Ikebe T, Chiba K, Shima T, Masuda C, Okuno R, Ohya H, Ogata K, Katsukawa C, Kawahara R, Tominaga K, Yabata J, Tada Y, Okabe N, Watanabe H, Chang B, Ogawa M, Ohnishi M, the Working group for beta-hemolytic streptococci in Japan. Evaluation of streptococcal toxic shock-like syndrome caused by group B streptococcus in adults

- in Japan between 2009 and 2013. *Journal of Infection and Chemotherapy*. 2015. 21 (3): 207-211.
- 14) Kohayagawa Y, Ishitobi N, Yamamori Y, Wakuri M, Sano C, Tominaga K, Ikebe T. Streptococcal toxic shock syndrome from necrotizing soft-tissue infection of the breast caused by a mucoid type strain. *Journal of Infection and Chemotherapy*. 2015. 21 (2): 144-147.
- 15) Okamoto F, Murakami K, Maeda E, Oishi A, Etoh Y, Kaida M, Makigusa M, Nakashima K, Jinnouchi Y, Takemoto H, Kakegawa H, Yamasaki C, Manabe S, Sasaki M, Ogata K, Ikebe T, Sera N. A Foodborne outbreak of group A streptococcal infection in Fukuoka prefecture, Japan. *Japanese Journal of Infectious Diseases*. 2014. 67 (4): 321-322.
- 16) Morimoto M, Tamura S, Hayakawa T, Yamanishi H, Nakamoto C, Nakamoto H, Ikebe T, Nakano1 Y, Fujimoto T. Phlegmonous gastritis associated with group A streptococcal toxic shock syndrome. *Intern Medicine*. 2014. 53 (22): 2639-2642.
- 17) Sanderson-Smith M, Oliveira D, Guglielmini J, McMillan D, Vu T, Holien J, Henningham A, Steer A, Bessen D, Dale J, Curtis N, Beall B, Walker M, Parker M, Carapetis J, Melderden L, Sriprakash K, Smeesters P, the M Protein Study Group. A systematic and functional classification of *Streptococcus pyogenes* that serves as a new tool for molecular typing and vaccine development. *Journal of Infectious Diseases*. 2014. 210 (8): 1325-1338.
- 18) Chang B, Wada A, Hosoya M, Oishi T, Ishiwada N, Oda M, Sato T, Terauchi Y, Okada K, Nishi N, Akeda H, Kamiya H, Ohnishi M, Ihara T, and the Japanese Invasive Disease Study Group. Characteristics of group B *Streptococcus* isolated from infants with invasive infections: A population-based study in Japan. *Japanese Journal of Infectious Diseases*. 2014. 67:356-360.
- 19) Oikawa J, Ishiwada N, Takahashi Y, Hishiki H, Nagasawa K, Takahashi S, Watanabe M, Chang B, and Kohno Y. Changes in nasopharyngeal carriage of *Streptococcus pneumoniae*, *Haemophilus influenzae* and *Moraxella catarrhalis* among healthy children attending a day-care centre before and after official financial support for the 7-valent pneumococcal conjugate vaccine and *H. influenzae* type b vaccine in Japan. *Journal of Infection and Chemotherapy*. 2014. 20:146-149.
- 21) Kasahara K, Komatsu Y, Koizumi A, Chang B, Ohnishi M, Muratani T, and Mikasa K. Serotype 35B *Streptococcus pneumoniae*, Japan, 2002-2012. *Journal of Infection and Chemotherapy*. 2014. 20:228-230.
- 22) Kuroki T, Ishida M, Suzuki M, Furukawa I, Ohya H, Watanabe Y, Konnai M, Aihara Y, Chang B, Ariyoshi K, Oishi K, Ohnishi M, and Morimoto K. Outbreak of *Streptococcus pneumoniae* serotype 3 pneumonia in extremely elderly people in a nursing home unit in Kanagawa, Japan, 2013. *Journal of the American Geriatrics Society*. 2014. 62:1197-1198.
- 24) Tamura K, Matsubara K, Ishiwada N, Nishi J, Ohnishi H, Suga S, Ihara T, Chang B, Akeda Y, Oishi K, the Japanese IPD Study Group. Hyporesponsiveness to the infecting serotype after vaccination of children with seven-valent pneumococcal conjugate vaccine following invasive pneumococcal disease. *Vaccine*. 2014. 32(42):1444-1450.
- 25) Ishiwada N, Hishiki H, Nagasawa K, Naito S, Sato Y, Chang B, Sasaki Y, Kimura K, Ohnishi M, and Shibayama K. The incidence of pediatric invasive *Haemophilus influenzae* and pneumococcal disease in Chiba prefecture, Japan before and after the introduction of conjugate vaccines. *Vaccine*. 2014. 32(42):5425-5231.
- 26) Akeda H, Chang B, Nakamura Y, Hamabata H, Ameku K, Toma T, Tamanaha E, and Ohnishi M. Impact of seven valent pneumococcal conjugate vaccine on nasopharyngeal carriage in young children in Okinawa, Japan. *World Journal of Vaccines*. 2015. 5(2):88-95.
- 27) Nishizuka M, Suzuki H, Ara T, Watanabe M, Morita M, Sato C, Tsuchida F, Seto J, Amemura-Maekawa J, Kura F, Takeda H.: A case of pneumonia caused by *Legionella pneumophila* serogroup 12 and treated successfully with imipenem. *J. Infect. Chemother*. 2014. 20:390-393.
- 28) Suzuki H, Nishizuka M, Ara T, Watanabe M, Morita M, Sato C, Tsuchida F, Seto J, Amemura-Maekawa J, Kura F, Takeda H.: Reply to the letter to the editor by Edelstein PH. *J. Infect.*

- Chemother. 2015. 1:77.
- 29) Koizumi N, Mizutani Muto M, Izumiya H, Suzuki M, Ohnishi M. Multiple-locus variable-number tandem repeat analysis and clinical characterization of *Leptospira interrogans* canine isolates. J Med Microbiol. 2105. 64(3): 288–294.
- 30) Kutsuna S, Kato Y, Koizumi N, Yamamoto K, Fujiya Y, Mawatari M, Takeshita N, Hayakawa K, Kanagawa S, Ohmagari N. Travel-related leptospirosis in Japan: a report on a series of five imported cases diagnosed at the National Center for Global Health and Medicine. J Infect Chemother. 2015. 21(3): 218–223.
- 31) Youn JH, Hayashida K, Koizumi N, Ohnishi M, Sugimoto C. Comparative genomic analysis of eight *Leptospira* strains from Japan and the Philippines revealing the existence of four putative novel genomic islands/islets in *L. interrogans* serovar Lai strain 56601. Comp Immunol Microbiol Infect Dis. 2014. 37(5–6): 289–297.
- 32) Toma C, Murray GL, Nohara T, Mizuyama M, Satou K, Koizumi N, Adler B, Suzuki T. Leptospiral outer membrane protein LMB216 is involved in enhancement of phagocytic uptake by macrophages. Cell Microbiol. 2014. 16(9): 1366–1377.
- 33) Yasuda SP, Gamage CD, Koizumi N, Nishio S, Isozumi R, Shimizu K, Koma T, Amada T, Suzuki H, Yoshimatsu K, Arikawa J. Distinct genetic characteristics of Sri Lankan *Rattus* and *Bandicota* (Murinae, Rodentia) inferred from mitochondrial and nuclear markers. Genes Genet Syst. 2014. 89(2): 71–80.
- 34) Shimizu Y, Sakamoto N, Ainoda Y, Hikone M, Kobayashi K, Iwabuchi S, Koizumi N, Ohnishi K. Leptospirosis in Japanese urban area: a case report and literature review. J Infect Chemother. 2104. 20 (4): 278–281.
- 35) Y. Takayama, S. Nakayama, K. Shimuta, T. Morita-Ishihara, and M. Ohnishi. Chracterization of azithromycin-resistant *Neisseria gonorrhoeae* isolated in Tokyo in 2005-2011. J. Infect. Chemother. 20:339-341. 2014.
- 36) Morita-Ishihara T, Unem M, Furubayashi K, Kawahata T, Shimuta K, Nakayama S, Ohnishi M. Treatment failure with 2 g of azithromycin (extended-release formulation) in gonorrhoea in Japan caused by the international multidrug-resistant ST1407 strain of *Neisseria gonorrhoeae*. Journal of Antimicrobial Chemotherapy. 69:2086-2090, 2014.
- 37) Nakao R, Kikushima K, Higuchi H, Obana N, Nomura N, Dongying B, Ohnishi M, Senpuku H. A Novel Approach for Purification and Selective Capture of Membrane Vesicles of the Periodontopathic Bacterium, *Porphyromonas gingivalis* : Membrane Vesicles Bind to Magnetic Beads Coated with Epoxy Groups in a Noncovalent, Species-Specific Manner. PLoS ONE. 2014. 9(5): e95137.
- 38) Tada A, Watanabe M, Senpuku H. Factors affecting changes in compliance with infection control practice by dentists in Japan. American Journal of Infection Control. 2015. 43(1): 95-97.
- 39) Bai D, Nakao R, Ito A, Uematsu H, Senpuku H. Immunoreactive antigens recognized in serum samples from mice intranasally immunized with *Porphyromonas gingivalis* outer membrane vesicles. FEMS Pathogens and Diseases. 2015. 73(3): 1-15.
- 40) Ohsumi T, Takenaka S, Wakamatsu R, Sakaue Y, Narisawa N, Senpuku H, Ohshima H, Terao Y, Okiji T. Residual structure of *Streptococcus mutans* biofilm following complete disinfection favors secondary bacterial adhesionand biofilm re-development. PLoS ONE. 2015. 10(1): e0116647.
- 41) Arai T, Kinoshita Y, and Senpuku H. Persistent colonization of *Candida albicans* yeast on the tongue in NOD/SCID.*e2f1<sup>-/-</sup>* mice. Journal of Infection and Chemotherapy. 2015, 21(5):370-375,
- 42) Arai T, Ochiai K, Senpuku H. *Actinomyces naeslundii* GroEL-dependent initial attachment and biofilm formation in a flow cell system. Journal of Microbiological Methods 2015. 109:160-166.
- 43) Golparian D, Shafer W, Ohnishi M, Unemo N. Importance of multi-drug efflux pumps in the antimicrobial resistance property of clinical multi-drug resistant isolates of *Neisseria gonorrhoeae*. Antimicrob Agents Chemother 2014, 58, 3556-3559.
- 44) Hayakawa K, Itoda I, Shimuta K, Takahashi H, Ohnishi M. A case of urethritis due to novel multilocus sequence type of the ST-11/ET-37 complex *Neisseria meningitidis* serogroup W in a MSM. Emerg Infect Dis 2014 20:1585-7.

- 45) Alam M, Rashed SM, Mannan SB, Islam T, Lizarraga-Partida ML, Delgado G, Morales T, Mendez JL, Navarro A, Watanabe H, Ohnishi M, Hasan NA, Huq A, Sack RB, Colwell RR, Cravioto A. Occurrence in Mexico, 1998 - 2008, of *Vibrio cholerae* CTX<sup>+</sup> El Tor Carrying an Additional Truncated CTX Prophage. Proc Natl Acad Sci U S A. 2014; 111: 9917-22.
- 46) Jacobsson S, Golparian D, Alm R, Huband M, Mueller J, Jensen J, Ohnishi M, Unemo M. High in-vitro activity of the novel spirocyclicimidinetrione AZD0914, a DNA gyrase inhibitor, against multidrug resistant *Neisseria gonorrhoeae* isolates suggests a new effective option for oral treatment of gonorrhea. Antimicrob Agents Chemother 2014; 58: 5585-5588.
- 47) Jacobsson S, Golparian D, Ly P, Ohnishi M, Hans F, Yat O, Magnus U. In vitro activities of the novel bicyclolides modithromycin (EDP-420, EP-013420, S-013420) and EDP-322 against multidrug-resistant clinical *Neisseria gonorrhoeae* isolates and international reference strains. J Antimicrobial Chemother 2015; 70: 173-177.
- 48) Igarashi T, Ito S, Sako M, Saitoh A, Hataya H, Mizuguchi M, Morishima T, Ohnishi K, Kawamura N, Kitayama H, Ashida A, Kaname S, Taneichi H, Tang J, Ohnishi M; Study group for establishing guidelines for the diagnosis and therapy of hemolytic uremic syndrome. Guidelines for the management and investigation of hemolytic uremic syndrome. Clin Exp Nephrol. 2014; 18: 525-557.
- 49) Muhammad I, Golparian D, Dillon JA, Johansson A, Ohnishi M, Sethi S, Chen SC, Nakayama SI, Sundqvist M, Bala M, Unemo M. Characterisation of blaTEM genes and types of beta-lactamase plasmids in *Neisseria gonorrhoeae* - the prevalent and conserved blaTEM-135 has not recently evolved and existed in the Toronto plasmid from the origin. BMC Infect Dis. 2014 Aug 22;14(1):454.
- 50) Lee K, Takano A, Taylor K, Sasika M, Shimozuru M, Konnai S, Kawabata H, Tsubota T. A Relapsing fever group *Borrelia* sp. similar to *Borrelia lonestari* found among wild sika deer (*Cervus nippon yesoensis*) and *Haemaphysalis* spp. ticks in Hokkaido, Japan. Ticks and Tick-borne Diseases. 2014; 5: 841-847.
- 51) Andoh M, Ogasawara Y, Sakata A, Ito T, Fujita H, Kawabata H, Ando S. Isolation of spotted fever group Rickettsia, *R. tamurae* and Candidatus *R. kotlanii*, from *Haemaphysalis megaspinosa* in Japan. Vector-Borne and Zoonotic Diseases. 2014; 14(9): 681-684.
- 52) Natsuaki M, Takada N, Kawabata H, Ando S, Yamanishi K. A case of tick-associated rash illness caused by *Amblyomma testudinarium*. Journal of Dermatology. 2014; 41(9): 834-836.
- 53) Takano A, Toyomane K, Konnai S, Ohashi K, Nakao M, Ito T, Andoh M, Maeda K, Watarai M, Sato K, Kawabata H. Tick surveillance for relapsing fever spirochete *Borrelia miyamotoi* in Hokkaido, Japan. PLoS One. 2014; 9(8): e104532.
- 54) Sato K, Takano A, Konnai S, Nakao M, Ito T, Kaneko M, Koyama K, Ohnishi M, Kawabata H. Human *Borrelia miyamotoi* infection in Japan. Emerging Infectious Diseases. 2014; 20(8): 1391-1393.
- 55) Konnai S, Hidano A, Yamada S, Gitahi, N, Higuchi H, Nagahata H, Ito T, Kawabata H, Ando S, Takano A, Murata S, Isesaki M. Suppressive effects of nutorophil by Salp16 Iper, a salivary gland protein, from *Ixodes persulcatus* Schulze tick. Insect Molecular Biology. 2014; 23(4): 466-474.
- 56) Takano A, Fujita H, Kadosaka T, Takahashi T, Yamauchi T, Ishiguro F, Takada N, Yano Y, Oikawa Y, Honda T, Gokuden M, Tsunoda T, Turumi M, Ando S, Andoh M, Sato K, Kawabata H. Construction of a DNA database for ticks collected in Japan: application of molecular identification based on the mitochondrial 16S rDNA gene. Medical Entomology and Zoology. 2014; 65(1): 13-21.
- 57) Suzuki S, Mimuro H, Kim M, Ogawa M, Ashida H, Toyotome T, Franchi L, Suzuki M, Sanada T, Suzuki T, Tsutsui H, Núñez G, Sasakawa C. Shigella IpaH7.8 E3 ubiquitin ligase targets glomulin and activates inflammasomes to demolish macrophages. Proc Natl Acad Sci U S A. 2014; 111:E4254-63.
- 58) Kiga, K., Mimuro, H., Suzuki, M., Shinozaki, A., Kobayashi, T., Sanada, T., Kim, M., Ogawa, M., Iwasaki, Y. W., Kayo, H., Fukuda-Yuzawa, Y., Yashiro, M., Fukayama, M., Fukao. T., and Sasakawa, C. Epigenetic silencing of miR-210 increases the proliferation of gastric epithelium during chronic Helicobacter pylori infection. Nat. Commun. 2014; 5:4497. doi: 10.1038/ncomms5497.
- 59) Harada-Hada, K., Harada, K., Kato, F., Hisatsune, J., Tanida, I., Ogawa, M., Asano, S., Sugai, M., Hirata M., Kanematsu. T. Phospholipase C-related

catalytically inactive protein participates in the autophagic elimination of *Staphylococcus aureus* infecting mouse embryonic fibroblasts. PLoS One 2014. 9: e98285.