

12. Department of Veterinary Science

- 1) Tsutsumi,R., Yoshida,Y., Suzuki,M., Imaoka,K., Yamamoto,O. Image Gallery: Annular erythema related to *Capnocytophaga canimorsus* bacteraemia after a dog bite. British Journal of Dermatology, 179(5):e196, 2018
- 2) Suzuki,M., Imaoka,K., Haga,Y., Mohri,M., Nogami,A., Shimojima,Y., Irie,Y., Sugiura,S., Morikawa,S. Characterization of three strains of *Capnocytophaga canis* isolated from patients with sepsis. Microbiology and Immunology, 62(9):567-573, 2018
- 3) Fukushima.K., Yanagisawa.N., Imaoka,K., Kimura,M., Imamura.A. Rat-bite fever due to *Streptobacillus notoymytis* isolated from a human specimen. The Journal of Infection and Chemotherapy, 24(4):302-304, 2018
- 4) Yumi Kumagai, Kozue Sato, Kyle R. Taylor, Aya Zamoto-Niikura, Koichi Imaoka, Shigeru Morikawa, Makoto Ohnishi, Hiroki Kawabata. A Relapsing fever group *Borrelia* sp. similar to *Borrelia theileri* is widely distributed among wild deer in Japan. Ticks and Tick-Borne Diseases, 9(3):465-470, 2018
- 5) Kei Yamamoto, Yasuyuki Kato, Yoshikazu Mutoh, Satoshi Kutsuna, Koichi Imaoka, Norio Ohmagari. Photo Quiz: A Traveler from Africa with Fever and Aggravated Chronic Back Pain. Clinical Infectious Diseases, 66(5):805-807, 2018
- 6) Okutani A, Inoue S, Morikawa S. Comparative genomics and phylogenetic analysis of *Bacillus anthracis* strains isolated from domestic animals in Japan. Infection, Genetics and Evolution. 2019. 71:128-139.
- 7) Shih, T.H., Chiang, J.T., Wu, H.Y., Tu, W.J., Inoue S, Tsai, C.T., Kuo, S.C., Yang, C.Y. and Fei, C.Y. (2018) An Analysis of Spillover Infection and Risk Factors for Human Exposure to Taiwan Ferret-badger Rabies in Taiwan. J. Prob. Health 6:1 doi:10.4172/2329-8901.1000196
- 8) Shiwa N, Kimitsuki K, Manalo DL, Inoue S, Park C-H. A pathological study of the tongues of rabid dogs in the Philippines. (2018) Arch Virol. doi:10.1007/s00705-018-3785-y.
- 9) Shih T-H, Chiang J-T, Wu H-Y, Inoue S, Tsai C-T, Kuo S-C, Yang C-Y, Fei C-Y. (2018) Human exposure to ferret badger rabies in Taiwan. Int. J. Environ. Res. Public Health. 15 1347; doi:10.3390/ijerph15071347.
- 10) Nguyen T.T., Nguyen V.D., Ngo C.G., Inoue S, Nguyen T.T.H., Pham N.T., Nguyen D.T., Nguyen H.D., and Nguyen T.K.A. (2018) Genetic characteristics of rabies virus cause endemic in Quang Nam and Quang Ngai, 2016-2017. Vietnam J. Prevent. Med. 28:102 – 110.
- 11) Shiwa N, Nakajima C., Kimitsuki K, Manalo D.L., Noguchi A., Inoue S, and Park C-H. (2018) Follicle sinus complexes (FSCs) in muzzle skin as postmortem diagnostic material of rabid dogs. J. Vet. Med. Sci. 80:1818-1821.
- 12) Inoue S, Park C-H. Hatta H. (2018) Production of monospecific polyclonal rabies virus antibodies in birds. Laboratory techniques in rabies. Fifth edition. Eds: Rupprecht CE., Fooks AR. and Abela-Ridder B. World Health Organization 2018.
- 13) Tian D, Uda A, Ami Y, Hotta A, Park ES, Nagata N, Iwata-Yoshikawa N, Yamada A, Hirayama K, Miura K, Koyama Y, Azaki M, Morikawa S. Protective effects of the Francisella tularensis ΔpdpC mutant against its virulent parental strain SCHU P9 in Cynomolgus macaques. Scientific reports, 2019, 9(1), 9193
- 14) Shimizu T, Otonari S, Suzuki J, Uda A, Watanabe K, Watarai M. Expression of Francisella pathogenicity island protein intracellular growth locus E (IgE) in mammalian cells is involved in intracellular trafficking, possibly through microtubule organizing center. MicrobiologyOpen, 2018, 8(4), e00684
- 15) Saha SS, Uda A, Watanabe K, Shimizu T, Watarai M. RtxA like protein contributes to infection of Francisella novicida in silkworm and human macrophage THP-1. Microbial pathogenesis, 2018, 123, 74-81.
- 16) Matsuno K, Nonoue N, Noda A, Kasajima N, Noguchi K, Takano A, Shimoda H, Orba Y, Muramatsu M, Sakoda Y, Takada A, Minami S, Une Y, Morikawa S, and Maeda K. Fatal Tickborne Phlebovirus Infection in Captive Cheetahs, Japan. Emerg. Infect. Dis. 2018, 24(9): 1726-1729.
- 17) Kimura T, Fukuma A, Shimojima M, Yamashita Y, Mizota F, Yamashita M, Otsuka Y, Kan M, Fukushi S, Tani H, Taniguchi S, Ogata M, Kurosu T, Morikawa S, Saijo M, Shinomiya H. Seroprevalence of severe fever with thrombocytopenia syndrome (SFTS) virus antibodies in humans and animals in Ehime prefecture, Japan, an endemic region of SFTS. J Infect Chemother. 2018, (10):802-806.
- 18) Saasa N, Kajihara M, Dautu G, Mori-Kajihara A, Fukushi S, Sinkala Y, Morikawa S, Mweene A, Takada A, Yoshimatsu K, Arikawa J. Expression of a Recombinant Nucleocapsid Protein of Rift Valley Fever Virus in Vero Cells as an Immunofluorescence Antigen and Its Use for Serosurveillance in Traditional Cattle Herds in Zambia. Vector Borne Zoonotic Dis. 2018, 18(5):273-277.
- 19) Yoshikawa T, Fujii H, Okutani A, Shibamura M, Omura N, Egawa K, Kato H, Inagaki T, Harada S, Yamada S, Morikawa S, Saijo M. Construction and characterization of bacterial artificial chromosomes harboring the full-length genome of a highly attenuated vaccinia virus LC16m8. PLoS One. 2018, 13(2): e0192725.
- 20) Zamoto-Niikura A, Tsuji M, Qiang W, Morikawa S, Hanaki KI, Holman PJ, Ishihara C. Babesia divergens Asia lineage is maintained between Ixodes persulcatus and sika deer in Hokkaido, Japan. Appl Environ Microbiol. 2018, pii: AEM.02491-17.
- 21) Tani H, Komeno T, Fukuma A, Fukushi S, Taniguchi S, Shimojima M, Uda A, Morikawa S, Nakajima N, Furuta Y, Saijo M. Therapeutic effects of favipiravir against severe fever with thrombocytopenia syndrome virus infection in a lethal mouse model: Dose-efficacy studies upon oral administration. PLoS One. 2018, 13(10):e0206416.