

16. Division of Experimental Animal Research

- 1) Zamoto-Niikura A, Tsuji M, Qiang W, Morikawa S, Hanaki KI, Holman PJ, Ishihara C. The *Babesia divergens* Asia Lineage Is Maintained through Enzootic Cycles between *Ixodes persulcatus* and Sika Deer in Hokkaido, Japan. Appl Environ. Microbiol. 84(7). pii: e02491-17, 2018
- 2) Kumagai Y, Sato K, Taylor KR, Zamoto-Niikura A, Imaoka K, Morikawa S, Ohnishi M, Kawabata H. A relapsing fever group *Borrelia* sp. is widely distributed among wild deer in Japan. Ticks Tick Borne Dis. 9:465–470, 2018
- 3) Fujii K, Sudaka Y, Takashino A, Kobayashi K, Kataoka C, Suzuki T, Iwata-Yoshikawa N, Kotani O, Ami Y, Shimizu H, Nagata N, Mizuta K, Matsuzaki Y, Koike S. VP1 Amino Acid Residue 145 of Enterovirus 71 Is a Key Residue for Its Receptor Attachment and Resistance to Neutralizing Antibody during Cynomolgus Monkey Infection. J Virol. 92: e00682-18, 2018
- 4) Li TC, Bai H, Yoshizaki S, Ami Y, Suzaki Y, Doan YH, Takahashi K, Mishiro S, Takeda N, Wakita T. Genotype 5 Hepatitis E Virus Produced by a Reverse Genetics System Has the Potential for Zoonotic Infection. Hepatol Commun. 3: 160–172, 2018
- 5) Hirai-Yuki A, Whitmire JK, Joyce M, Tyrrell DL, Lemon SM. Murine models of hepatitis A virus infection. Cold Spring Harb Perspect Med. 9:a031674, 2019